



Synthesis of 1-benzyloxy pyrazin-2(1H)-one derivatives



Anh Hung Mai, Sonalika Pawar[†], Wim M. De Borggraeve^{*}

Molecular Design and Synthesis, Chemistry Department, University of Leuven, Celestijnenlaan 200F-box 2404, 3001 Heverlee, Belgium

ARTICLE INFO

Article history:

Received 30 May 2014

Revised 23 June 2014

Accepted 25 June 2014

Available online 1 July 2014

Keywords:

Aspergillilic acid

Hydroxamic acid

Pyrazin-2(1H)-one

Amidation

ABSTRACT

Different approaches for the synthesis of 1-benzyloxy pyrazin-2(1H)-one derivatives from simple amino acids have been investigated. A library of 33 precursors for the preparation of *N*-hydroxy pyrazinones was obtained in moderate to good yields.

© 2014 Elsevier Ltd. All rights reserved.

The pyrazinone is a valuable scaffold in medicinal chemistry. It is present amongst others in the inhibition of HCV NS3 protease,^{1,2} neutrophil elastase,³ prolyl oligopeptidase,⁴ TF-FVIIa⁵ and thrombin.^{6,7} A lot of research in our group was directed towards the development of synthetic strategies for highly functionalized pyrazinones.^{8–15} Since there are bioactive compounds known in nature containing an *N*-hydroxypyrazinone core (e.g., aspergillilic acid, Fig. 1),^{16–18} we focused our attention to the development of new strategies to synthesize pyrazinones containing 1-benzyloxy functionality. These compounds can potentially give rise to aspergillilic acid-like hydroxamic acids upon deprotection. Furthermore, in extension of our work on 1-alkyl/aryl functionalized pyrazinone-3-carboxamides,¹⁵ we also intended to prepare the corresponding 1-benzyloxy-3-carboxamide pyrazinones.

Our synthetic scheme for the synthesis of 1-benzyloxy pyrazin-2(1H)-ones improves upon reported procedures^{19–23} and is similar to chemistry we already applied in the preparation of 1-alkyl/aryl pyrazinones.^{11,12} It relies on the base catalysed condensation of a glyoxal derivative with an amino acid hydroxamate. The condensations with phenyl glyoxal and diacetyl need a higher reaction temperature (50–70 °C) compared to those with glyoxal and methyl glyoxal. In order to avoid excessive side reactions resulting in very complex mixtures, it was important to use the glyoxal derivative as a limiting reagent (0.9 equiv, see Supplementary information) and to add it slowly to the reaction mixture via a syringe pump over the

course of 30–120 min. The regioselectivity of this reaction with an unsymmetrical glyoxal derivative (methyl/phenyl glyoxal) was reported before^{19,20} and was confirmed by us via NMR analysis. Proton H-6 in compounds **3b,c,e,f,m,n,q,t,w,y,zc,zg** (unsubstituted at position 6, but substituted at position 5) shows a clear NOE correlation with the methylene protons of the *O*-benzyl group, as well as an HMBC correlation with C-2 (Fig. 2).

The synthesis of 1-benzyloxy-3-alkylpyrazin-2(1H)-ones (**3a–f**, Scheme 1) was achieved by amidation of *N*-Boc protected amino acids **1** using *O*-benzyl hydroxylamine in combination with HOBt/EDCI/DIPEA/DMF, followed by Boc-deprotection and condensation. *O*-Benzyl hydroxamate **2a** could also be prepared via direct




Figure 1. Aspergillilic acid (**1**).





Figure 2. NOE and HMBC correlations of H-6 in compounds **3**.

^{*} Corresponding author. Tel.: +32 16 327693; fax: +32 16 327990.

E-mail address: wim.deborggraeve@chem.kuleuven.be (W.M. De Borggraeve).

[†] Current address: Chemistry Department, Fergusson College, F.C. Road, 411004 Pune, India.



Scheme 1. Synthetic pathway for preparation of 1-benzoyloxy-3-alkylpyrazin-2(1H)-ones. Reagents and conditions: (a) BnONH₂·HCl (1 equiv), HOBT (1.3 equiv), EDCI (1.3 equiv), DIPEA (2.3 equiv), DMF, −10 °C then rt, 16 h; (b) (i) 4 M HCl (10 equiv) in dioxane, rt, 30 min. (ii) R⁵R⁶(CO)₂ (0.9 equiv), 2 M NaOH, pH 8–10, MeOH–H₂O (2:1), −35 °C then rt, overnight; (c) BnONH₂ (1.1 equiv), LiHMDS (3.1 equiv), THF, −78 °C, 2 h.


amidation of the methyl ester of amino acid **4** using LiHMDS base in THF.²⁴ A slight reduction in yield and a shorter reaction time are observed in this instance (**Scheme 1**, entry **2a**). The direct amidation route could also be applied to more complex substrates as exemplified below in **Scheme 3**. Using this approach of slow addition with glyoxal as the limiting reagent, the previously reported 1-benzoyloxy-3-alkylpyrazin-2(1H)-ones **3a**, **3b** and **3d** were obtained in better yields as compared to the literature.^{21–23}

The synthesis of the novel and more complex 1-benzoyloxy-3-carboxamides **3g–zg** is described in **Schemes 2 and 3**. In this pathway, the amino group in diethyl amino malonate ester hydrochloride (**5**) is first Boc-protected to form **6**. This is followed by iterative mono-saponification amidation to generate **10** (**Scheme 2**). These compounds are then converted to pyrazine-2(1H)-ones in moderate to good yields after Boc-removal (**Schemes 2 and 3**). Compounds **10** can alternatively directly be obtained via conversion of ethyl ester **11** using LiHMDS and NH₂OBn (**Scheme 3**). In the case of 3-carboxylated derivatives of 1-benzoyloxy-3-alkylpyrazin-2(1H)-one, the latter approach is able to


reduce the number of reaction steps leading to the final products; however, it does limit the late stage diversification at C-3 of the target compounds, which in terms of library generation is a drawback.

In a further effort to shorten the synthetic protocol in the synthesis of 1-benzoyloxy-3-alkylpyrazin-2(1H)-ones **3**, we performed the cyclization of the product of Boc-deprotection of **8** with glyoxals to generate 1-benzoyloxy-3-alkylpyrazin-2(1H)-one 3-carboxyl ethyl esters **12**, which could be used as precursors in a one-step amidation to form **3** (**Scheme 4**) using MgCl₂ as Lewis acid catalyst.²⁵ The desired secondary amide products (**3zc**, **3zd** and **3zg**) could be obtained in high yields by treating ethyl esters **12** with primary amines. However, no conversion was detected in case of secondary amines even after prolonged reaction time and heating at a temperature of 50 °C (**3zh** and **3p** however can be synthesized via methods described in **Schemes 2 and 3**).

Using these approaches, a 33-component library of 1-benzoyloxy-3-alkylpyrazin-2(1H)-one derivatives, precursors for the synthesis of N-hydroxypyrazinones, has been prepared in moderate to good



Scheme 2. Synthetic pathway for preparation of 1-benzoyloxy-3-carboxamides **3**. Reagents and conditions: (a) Boc₂O (1.05 equiv), NaHCO₃ (1.05 equiv), DMAP (0.01 equiv), H₂O–dioxane, rt, overnight; (b) KOH (1 equiv), EtOH, rt, overnight; (c) BnONH₂·HCl (1 equiv), HOBT (1.3 equiv), EDCI (1.3 equiv), DIPEA (2.3 equiv), DMF, −10 °C then rt, 16 h; (d) R⁷R⁸NH (1 equiv), HOBT (1.3 equiv), EDCI (1.3 equiv), DIPEA (1.3 equiv), DMF, −10 °C then rt, 16 h; (e) (i) 4 M HCl (16 equiv) in dioxane, rt, 30 min; (ii) R⁵R⁶(CO)₂ (0.9 equiv), 2 M NaOH, pH 8–10, MeOH–H₂O (2:1), −35 °C then rt, overnight.



Compd.	R ⁵	R ⁶	R ⁷	R ⁸	Yield (%)
11a	-	-	Et	Et	84
11b	-	-	H	3-FC ₆ H ₄ CH ₂	73
11c	-	-	H	4-FC ₆ H ₄ CH ₂	80
11d	-	-	H	4-ClC ₆ H ₄ CH ₂	69
11e	-	-	H	3,4-Cl ₂ C ₆ H ₃ CH ₂	78
10a	-	-	Et	Et	79
10i	-	-	H	3-FC ₆ H ₄ CH ₂	66
10j	-	-	H	4-FC ₆ H ₄ CH ₂	64
10k	-	-	H	4-ClC ₆ H ₄ CH ₂	57
10l	-	-	H	3,4-Cl ₂ C ₆ H ₃ CH ₂	67
3y	Me	H	Et	Et	42
3z	H	H	H	3-FC ₆ H ₄ CH ₂	51
3za	Me	Me	H	3-FC ₆ H ₄ CH ₂	47
3zb	H	H	H	4-FC ₆ H ₄ CH ₂	60
3zc	Me	H	H	4-FC ₆ H ₄ CH ₂	58
3zd	H	H	H	4-ClC ₆ H ₄ CH ₂	67
3ze	Me	Me	H	4-ClC ₆ H ₄ CH ₂	42
3zf	H	H	H	3,4-Cl ₂ C ₆ H ₃ CH ₂	61
3zg	Me	H	H	3,4-Cl ₂ C ₆ H ₃ CH ₂	49

Scheme 3. Direct amidation of ethyl ester **11**. Reagents and conditions: (a) R⁷R⁸NH (1 equiv), HOBT (1.3 equiv), EDCI (1.3 equiv), DIPEA (1.3 equiv), DMF, -10 °C then rt, 16 h; (b) BnONH₂ (1.1 equiv), LiHMDS (4.1 equiv), THF, -78 °C, 2 h; (c) (i) 4 M HCl (16 equiv) in dioxane, rt, 30 min. (ii) R⁵R⁶(CO)₂ (0.9 equiv), 2 M NaOH, pH 8–10, MeOH–H₂O (2:1), -35 °C then rt, overnight.

12

Compd.	R ⁵	R ⁶	R ⁷	R ⁸	Yield (%)
12a	H	H	-	-	68
12b	Me	H	-	-	61
3zc	Me	H	H	4-FC ₆ H ₄ CH ₂	95
3zd	H	H	H	4-ClC ₆ H ₄ CH ₂	90
3zg	Me	H	H	3,4-Cl ₂ C ₆ H ₃ CH ₂	90
3zh	H	H	Me	Bn	0 ^(c)
3p	H	H	Me	4-FC ₆ H ₄ CH ₂	0 ^(c,d)

Scheme 4. Synthesis of **3** via amidation of ester **12**. Reagents and conditions: (a) (i) 4 M HCl (16 equiv) in dioxane, rt, 30 min. (ii) R⁵R⁶(CO)₂ (0.9 equiv), 2 M NaOH, pH 7–8, MeOH–H₂O (2:1), -35 °C then rt, overnight; (b) (i) MgCl₂ (2 equiv), THF, rt, 5 min. (ii) R⁷R⁸NH (2.5 equiv), rt, 16 h; (c) No conversion, **12a** was recovered (by LC–MS); (d) 38% yield of **3p** (from **8**, Scheme 2).

yields with minimal reaction steps. Pyrazin-2(1H)-ones **3c**, **3e**, **3f** and 27 other 3-carboxamide substituted analogues **3g–zg** are new compounds. Deprotection of this library will be subject of another paper.

Acknowledgment

The authors gratefully acknowledge financial support from KU Leuven via OT/11/047 and the Ministry of Education and Training (Vietnam International Education Development, VIED).

Supplementary data

Supplementary data (general procedures for synthesis and characterization of all compounds, copies of ¹H and ¹³C NMR spectra of compounds **3** and **12**) associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.tetlet.2014.06.100>. These data include MOL files and InChIKeys of the most important compounds described in this article.

References and notes

- Örtqvist, P.; Gising, J.; Ehrenberg, A. E.; Vema, A.; Borg, A.; Karlén, A.; Larhed, M.; Danielson, U. H.; Sandström, A. *Bioorg. Med. Chem.* **2010**, *18*, 6512–6525.
- Gising, J.; Belfrage, A. K.; Alogheli, H.; Ehrenberg, A.; Åkerblom, E.; Svensson, R.; Artursson, P.; Karlén, A.; Danielson, U. H.; Larhed, M.; Sandström, A. *J. Med. Chem.* **2013**, *57*, 1790–1801.
- Hansen, P.; Ivarsson, M.; Lawitz, K.; Lönn, H.; Nikitidis, A.; Ray, A. U.S. Patent 8,114,881, Feb. 14, 2012.
- Haffner, C. D.; Diaz, C. J.; Miller, A. B.; Reid, R. A.; Madauss, K. P.; Hassell, A.; Hanlon, M. H.; Porter, D. J. T.; Becherer, J. D.; Carter, L. H. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 4360–4363.
- Zhang, X.; Glunz, P. W.; Jiang, W.; Schmitt, A.; Newman, M.; Barbera, F. A.; Bozarth, J. M.; Rendina, A. R.; Wei, A.; Wen, X.; Rossi, K. A.; Luetgten, J. M.; Wong, P. C.; Knabb, R. M.; Wexler, R. R.; Scott Priestley, E. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 1604–1607.
- Young, M. B.; Barrow, J. C.; Glass, K. L.; Lundell, G. F.; Newton, C. L.; Pellicore, J. M.; Rittle, K. E.; Selnick, H. G.; Stauffer, K. J.; Vacca, J. P.; Williams, P. D.; Bohn, D.; Clayton, F. C.; Cook, J. J.; Krueger, J. A.; Kuo, L. C.; Lewis, S. D.; Lucas, B. J.; McMasters, D. R.; Miller-Stein, C.; Pietrak, B. L.; Wallace, A. A.; White, R. B.; Wong, B.; Yan, Y.; Nantermet, P. G. *J. Med. Chem.* **2004**, *47*, 2995–3008.
- Kranjc, A.; Mašić, L. P.; Reven, S.; Mikic, K.; Preželj, A.; Stegnar, M.; Kikelj, D. *Eur. J. Med. Chem.* **2005**, *40*, 782–791.
- Vekemans, J.; Pollers-Wieërs, C.; Hoornaert, G. *J. Heterocycl. Chem.* **1983**, *20*, 919–923.
- De Borggraeve, W. M.; Verbist, B. M. P.; Rombouts, F. J. R.; Pawar, V. G.; Smets, W. J.; Kamoun, L.; Alen, J.; Van der Eycken, E. V.; Compennolle, F.; Hoornaert, G. *J. Tetrahedron* **2004**, *60*, 11597–11612.
- Heeres, J.; de Jonge, M. R.; Koymans, L. M. H.; Daeyaert, F. F. D.; Vinkers, M.; Van Aken, K. J. A.; Arnold, E.; Das, K.; Kilonda, A.; Hoornaert, G. J.; Compennolle, F.; Cegla, M.; Azzam, R. A.; Andries, K.; de Béthune, M.-P.; Azijn, H.; Pauwels, R.; Lewi, P. J.; Janssen, P. A. J. *J. Med. Chem.* **2004**, *48*, 1910–1918.
- Azzam, R.; De Borggraeve, W.; Compennolle, F.; Hoornaert, G. *J. Tetrahedron Lett.* **2004**, *45*, 1885–1888.
- Azzam, R.; De Borggraeve, W. M.; Compennolle, F.; Hoornaert, G. *J. Tetrahedron* **2005**, *61*, 3953–3962.
- Pawar, V. G.; De Borggraeve, W. M. *Synthesis* **2006**, 2799–2814.
- Kamoun, L.; De Borggraeve, W. M.; Gielen, C.; Voet, A.; Robeyns, K.; De Maeyer, M.; Van Meervelt, L.; Compennolle, F.; Hoornaert, G. *Eur. J. Org. Chem.* **2007**, 2977–2986.
- Pawar, S. V.; Pawar, V. G.; Dehaen, W.; De Borggraeve, W. M. *Org. Lett.* **2008**, *10*, 4473–4476.
- Dutcher, J. D. *J. Biol. Chem.* **1947**, *171*, 321–339.
- Lott, W. A.; Shaw, E. *J. Am. Chem. Soc.* **1949**, *71*, 70–73.
- Hogg, R. W.; Biswas, C. S.; Broquist, H. P. *J. Bacteriol.* **1965**, *90*, 1265–1270.
- Dunn, G.; Ramsay, D. W. C.; Elvidge, J. A.; Spring, F. S.; Newbold, G. T.; Sweeny, W. *Nature* **1949**, *164*, 181.
- Jones, R. G. *J. Am. Chem. Soc.* **1949**, *71*, 78–81.
- Katoh, A.; Ohkanda, J.; Itoh, Y.; Mitsuhashi, K. *Chem. Lett.* **1992**, *21*, 2009–2012.
- Ohkanda, J.; Tokumitsu, T.; Mitsuhashi, K.; Katoh, A. *Bull. Chem. Soc. Jpn.* **1993**, *66*, 841–847.
- Ohkanda, J.; Kumasaka, T.; Takasu, A.; Hasegawa, T.; Katoh, A. *Heterocycles* **1996**, *43*, 883–889.
- Gissot, A.; Volonterio, A.; Zanda, M. *J. Org. Chem.* **2005**, *70*, 6925–6928.
- Guo, Z.; Dowdy, E. D.; Li, W.-S.; Polniaszek, R.; Delaney, E. *Tetrahedron Lett.* **2001**, *42*, 1843–1845.